



In This Issue

- Upcoming Meetings
- Results: Statewide 2017 Small Grain Variety Trials
- Southern Blight Cliff Notes - 2017
- Destroying weed seed at harvest: seed destroying technology to be tested in the Sacramento Valley

Submitted by:

Sarah Light
UCCE Farm Advisor
Sutter-Yuba and
Colusa Counties

Two Great Events Coming Up!

Soils of the Southern Sacramento Valley Field Tour

Thursday, October 12, 2017 8 a.m. - 12.15 p.m.

Rominger Brothers' and Triad Farms

3.0 CCA Credits Pending

Join us for a half-day workshop profiling some of the major soils of the southern Sacramento Valley. We will be discussing several decision making tools related to irrigation management and nutrient toxicity/ deficiency issues associated with local soils. Coffee and donuts provided. See [Sacramento Valley Field Crops Website](#) for additional agenda details.



It's Alive!

Principles and Practices to Build Soil Health on Your Farm

Thursday, October 12, 2017 8:30 a.m. – 12 noon

UC Davis Russell Ranch Sustainable Agriculture Facility,
Kinsella Lane, Winters, CA

This workshop is for all farmers who want to learn about soil health. It will focus on organic systems, but will include principles and practices relevant to all systems. Through demonstrations, hands-on exercises and field observations, participants will learn:

- How the living part of the soil affects important soil properties
- How to assess your farm's soil health using your 5 senses
- How soil management practices affect soil health

Advance registration required. To register, please visit <http://asi.ucdavis.edu/event-calendar/soilhealth>

ANNOUNCEMENT!

The results of the Statewide 2017 Small Grain Variety Trials are in! Mark Lundy, Statewide Specialist in Grain Cropping Systems, has published the results from this year's variety trials. Check them out here:

http://smallgrains.ucanr.edu/Variety_Results/2017/

Southern Blight Cliff Notes- 2017

*Cassandra Swett, Assist Cooperative Extension Specialist, ANR
Department of Plant Pathology, University of California, Davis
Joe Nunez, UCCE Farm Advisor, Emeritus, Kern County*

This was a big year for southern blight. Out of over forty crown rot samples received in the UC Davis Vegetable and Agronomic Crop Pathology lab (Swett lab) in summer 2017, 50% were southern blight, 40% were Fusarium root and crown rot, and 10% were attributed to other causes. This disease was commonly reported to cause over 50% mortality in affected fields.

Southern blight is a very destructive, fast acting crown rot disease that rapidly kills the plant. Over 500 different plants are southern blight hosts. Affected crops that came through the Swett lab included pepper, potato, tomato, cucumber, canary bean, chard, and sunflower. Most unusually, this disease caused major losses in many northern counties in the San Joaquin and Sacramento valley, where it is not typically an issue, including Yolo, Colusa, Contra Costa, San Joaquin, and Merced.

Southern blight is not typically considered to be a widespread problem in California--major impacts are usually restricted to the Kern County area; the widespread distribution we saw this year is NOT likely due to pathogen spread to new fields. Southern blight is favored by high temperatures (over 86°F), high soil moisture, dense canopies, and frequent irrigation. It seems most plausible that a combination of late planting dates and record high summer temperatures created unusually favorable conditions for the pathogen in the northern part of the valley.

Although not a new disease to the state, the increased damage from the disease this year may mean that this will be a bigger issue next year, if the environment is conducive and the disease is not properly managed. Southern blight is caused by the fungus *Sclerotium rolfsii*. The fungus survives in soil as hardened structures called sclerotia for at least five years. Each infected plant can literally produce tens of thousands of sclerotia and then become more widely distributed in a field with each successive field operation. Although initially this disease may only affect a few plants in the field, southern blight can be serious enough to cause significant yield loss within a season or two. With a host range of over 500 plants, this fungus can easily persist from year to year in infected crop debris.

How to identify southern blight in the field

Southern blight misdiagnosis is likely if it occurs in an area where it has not historically been an issue. Scouting and mapping infested locations in fields during the summer months will greatly help in determining what options can be taken before the sclerotia levels become too numerous and cause severe crop loss. It can be easy to confuse southern blight with other crown rotting diseases, for example Fusarium crown rot. Accurate diagnosis is critical to effective control. You can distinguish southern blight based on the following diagnostic traits, one or more of which is often, but not always present. Part of the trick to diagnosis is not to just look at the plant, but also look at the soil right around the crown.


These small, tan to reddish brown sclerotia form at the base of the plant and / or in the soil right around the plant. The sclerotia look like alfalfa seeds when young but turn brown with age. (Photo credit: J. Nunez)



White fungal mycelium (thread-like strands) growing INTO the soil. No other fungus will grow extensively in the soil. Sometimes you also see sclerotia in the soil. (Photo credit: R. C. Swett, L: J. Nunez)



White fan like mycelial (thread like) grown on the crown / affected tissues. Severely affected plants can have vascular discoloration, which may be confused with Fusarium wilt. (Photo credit: J. Nunez)



Plants go from looking healthy to dead in less than a week—this is much faster than most crown rots. (Photo credit: C. Swett)



In affected fields, the disease patches are roughly circular. From a distance, they look like bands of dead plants. (Photo credit: L: J. Nunez, R: C. Swett)



If none of these characteristics are present, the best way to diagnose the disease is to put infected tissue in a plastic bag on a moist paper towel and leave it at room temperature. The southern blight fungus will produce a distinct fan like growth within about 5-7 days. After about 5-14 days it will make round white balls that then turn into amber colored sclerotia. Both the fan growth and the sclerotia are unique to this fungus. (Photo credit: C. Swett)



In-season Fungicide Applications

Southern blight acts fast, so as soon as you detect the problem, it's critical to get out there to spray. Fungicides work by covering the crown tissue both above and below the soil, killing the fungus around the crown. For vegetable crops, fungicides such as flutolanil, penthiopyrad, and tebuconazole are known to be effective in the management of southern blight. However, these products are registered on only a few vegetables so make sure to check crop registration before using these on any vegetable crop. Also, some of these fungicides have severe plant-back restrictions so crop rotations need to be carefully planned. As always make sure to read and follow label directions to avoid any problems.

Perhaps the biggest obstacle to fungicide control of southern blight is application timing and method. Because southern blight is basically a summer time disease it rears its ugly head when most crops are near maturity with a full canopy cover. Getting fungicides to the base of the stem and onto the surface of the soil is very difficult especially for fields on drip irrigation systems. Chemigation through sprinklers is a better option especially on crops like garlic and onions which do not have a dense canopy.

If the crop has a dense canopy that the fungicide cannot penetrate, then a fungicide application won't work to control the disease. Good, narrow canopy crops for fungicide control include onions, garlic, beans, sunflower, and potatoes to a certain extent. Poor, dense canopy crops for fungicide control include tomatoes, melons, peppers, potatoes, and vegetables grown for seed, such as lettuce.

I received the question several times on whether it's possible to apply the fungicide by drip chemigation (through the drip line). If it's buried drip, then no—it would take a lot of water to get to the soil surface, which would be likely to cause other problems; several trials have been conducted in the past in Kern County, and it's never worked. Surface drip would work, but this irrigation method is not common in California.

Managing soil moisture

Manipulating your irrigation to maintain a dry surface may help reduce losses if you detect the fungus in your field. The one advantage of drip irrigation is that the soil surface can more easily be kept dry which inhibits infection by *Sclerotium rolfsii*. However, alternating wet and dry periods can be a problem--wet periods followed by dry episodes can be particularly conducive to disease development.

Crop rotation

If you have detected southern blight in your field, one of the best things you can do the following year is to plant a crop that you can effectively fungicide treat in-season, such as onion, garlic, or any crop that is irrigated on sprinklers. The disease can be effectively controlled in these crops, preventing sclerotia from increasing.

Crop rotations with non-host crops are limited because of the wide host range. Poor-host crops such as corn and small grains (wheat, millet, oats) can help to significantly reduce sclerotia levels in the field. Most, if not all, of these crops can become infected by the fungus, but either they are not good hosts and/or the environmental conditions during the growing season are not conducive to pathogen growth. For instance, wheat can host the fungus, but it's too cold for fungal growth during the time that wheat is grown. On the other hand, rotation with legumes such as beans, peas and hairy vetch will greatly increase soil infestation

levels. Mustard cover crops can suppress southern blight, and may be useful for organic producers, where fumigation is not an option.

Soil treatment

Once sclerotia levels become too numerous in a field then fumigation will need to be considered. Fumigation with metam sodium (Vapam, K-Pam) can be effective, but ideally it needs to be applied through sprinklers so it percolates down into the soil at least 6" to kill the fungus in the soil zone where it is active. Because of restrictions in application, sprinkler application is not allowed in many counties, so you have to shank it in. This method of application is not as effective since the fumigant does not penetrate deep enough into the soil. Fields with shanked applications may still suffer major southern blight losses. Also, the requirement of buffer zones for metam applications means the field may become re-infested in short time as sclerotia are moved from the buffer zones into the rest of the field with various tractor operations.

Deep plowing will bury the sclerotia and get it away from attacking plants at the soil line. Sclerotia deeper than 6 inches are usually parasitized by other microbes and are killed over time. Of course plowing is not an option for fields where buried drip irrigation systems are already installed.

Sclerotia near the surface of the soil can be killed when exposed to high temperatures (105-120°F) for two to four weeks during the summer months. Solarization alone is not generally considered a viable management strategy, but when soils were solarized before addition of a biological control or a fungicide, disease was reduced by 70-100% compared to the same biological or chemical treatment without solarization. Cover and secure clear polyethylene plastic over moistened soil. Make sure to prepare the soil for planting before solarizing, since cultivation and the incorporation of amendments can bring buried sclerotia back to the upper soil layers.

There are several fungi that appear to have some antagonistic effects on southern Blight including RootShield (*Trichoderma harzianum*). There are no field studies that indicate efficacy of bacterial products (eg. Serenade Soil) and to the authors' knowledge there are no studies to support the use of plant defense inducing products such as Regalia.

Disease resistance

For most crops, southern blight resistant cultivars are not available. However, for vegetable crops such as tomatoes there are some rootstocks reported to be resistant to southern blight, which are currently under study in field trials in California. These may be a promising option for small scale and organic producers.

Crystal Gazing--What's going to happen next year?

This was an unusually hot summer and crops went in late due to late spring rains—this combination of factors likely accounts for the widespread occurrence. If crops are planted on time next year, and / or it's not so hot, then the disease might not rear its ugly head. But one thing to keep in mind for folks that have fields with southern blight this year—now, inoculum levels are higher, so it's going to take less to make this a problem next year.

Destroying weed seed at harvest: seed destroying technology to be tested in the Sacramento Valley

Mariano Galla— UCCE Agronomy and Weed Science Advisor-- Glenn, Butte, and Tehama Counties

Weed control in field crops is a major concern for growers all around the world. In the absence of an adequate weed control program, weed pressure can reduce yields by as much as 48%. Almost all weed management systems rely on the use of herbicides, resulting in widespread development of herbicide-resistant weeds that pose a serious threat to crop productivity worldwide. New approaches to weed control management that include both chemical and non-chemical tools are, thus, necessary for the development of sustainable weed control systems.

Currently, in field crops, standard weed management programs target weeds either at germination (using pre-emergent herbicides) or at early seedling stage. The overall goal is to keep the field weed-free early in the growing season, when crops are particularly vulnerable to weed competition. However, inevitably some weeds are able to escape herbicide control and survive to maturity producing significant amounts of seeds that replenish the seedbank in the soil.

In Australia, because of the rapid spread of resistance to multiple herbicides in annual ryegrass and wild radish, researchers have increased the effort in developing new control tools. Studies found that the vast majority of in-crop annual ryegrass germinates from the previous year's seed production. Furthermore, up to 80% of the total ryegrass seeds are collected by the combine during harvest and returned to the field in the chaff fraction.

Thus, Australian researchers have tried to minimize additions to the seed bank through developing harvest weed seed control systems (HWSC). Many different systems have been tested such as chaff carts that collect the chaff exiting the combine, windrow burning, and the direct-bale system.

Based on the intuition of Ray Harrington, an Australian grower, researchers developed the Harrington Seed Destructor (HSD): a trailer-mounted cage mill with chaff and straw transfer system powered by a diesel motor. Chaff exiting the combine is delivered to the cage mill where it is crushed before being returned to the field (figure 1). In field trials the HSD was able to destroy at least 95% of ryegrass, wild oat and wild radish seeds present in the chaff fraction of the harvest residue. The HSD was commercialized in 2013 and in 2016 an integrated system that can be mounted within the rear of the combine was developed (figure 2).

The HSD can be a useful tool in an integrated weed management system that includes herbicide applications early in the season. In Australia, it has been reported that the combination of herbicide early in the growing season with HWSC successfully reduced annual ryegrass population pressure over the years.

To date, HWSC are being tested across the US in different locations. De Bruin engineering, the company that commercialized the integrated Harrington Seed Destructor in Australia, is interested in running a demonstration trial in the Sacramento Valley. If you are interested in the Harrington Seed Destructor please reach out to Mariano Galla (mfgalla@ucanr.edu) in the Orland UCCE office, Konrad Mathesius (kpmathesius@ucanr.edu) in the Woodland UCCE office or Sarah Light (selight@ucanr.edu) in the Yuba City UCCE office.

Walsh, Michael, Peter Newman, and Stephen Powles. "Targeting weed seeds in-crop: a new weed control paradigm for global agriculture." Weed Technology 27.3 (2013): 431-436.




Figure 1 Schematic of Harrington Seed Destructor. Source Wash et al 2012



Figure 2 Ray Harrington and his integrated Harrington Seed Destructor. Source: <http://www.abc.net.au/news/2017-06-15/ray-harrington-and-tractor-with-seed-destructor/8620674>

ANR NONDISCRIMINATION AND AFFIRMATIVE ACTION POLICY STATEMENT FOR UNIVERSITY OF CALIFORNIA.

May, 2015 It is the policy of the University of California (UC) and the UC Division of Agriculture & Natural Resources not to engage in discrimination against or harassment of any person in any of its programs or activities (Complete nondiscrimination policy statement can be found at <http://ucanr.edu/sites/anrstaff/files/215244.pdf>). Inquiries regarding ANR's nondiscrimination policies may be directed to John I. Sims, Affirmative Action Compliance Officer/Title IX Officer, University of California, Agriculture and Natural Resources, 2801 Second Street, Davis, CA 95618, (530) 750-1397.