



“OLD” VS “NEW” FIELDS: EXPLORING CAUSAL FACTORS FOR THE YIELD GAP

Patricia Lazicki
UCCE Vegetable Crops Advisor
Yolo, Solano & Sacramento counties
palazicki@ucanr.edu; 530-219-5198

EXPERIMENTAL DESIGN




- 3 field pairs (old vs new (from walnut)), 5 locations/field
- Match:
 - Soil type
 - Variety
 - Approximate planting time (Late March/ early April)
- Measure biological, physical, chemical indicators


Location	Soil type	Old/New	Variety	Harvest date
Madison	Tehama loam (light surface soil, clay layer)	Old	SVTM 9027	8-Aug
		New	SVTM 9027	4-Aug
Winters	Brentwood silty clay loam (heavier soil, clay layer)	Old	HM 58841	18-Aug
		New	HM 58841	31-Jul
Zamora	Yolo silt loam (medium texture, no subsurface layer)	Old	HM 0371	1-Aug, 12-Aug
		New	HM 0371	30-Aug

Factor Type	Analyses	Timing (1 st year)
Biological	Greenhouse assays; sterilized vs unsterilized soils	Planting
	Nematode communities	Planting
	Microbial community structure (PLFA)	Planting
	Declined plants, pathogen identities	Pre-harvest
Microbial communities (DNA)		Planting
Physical	Bulk density (0-6", 6-12", 12-18", 18-24")	Early season
	Aggregate size distribution	Planting
	Moisture content (center vs edge, 0-24")	Midseason
	Soil pit, root distribution (observational, only)	Mid - late season
Chemical	Basic fertility	Planting
	Nutrient distributions (center vs edge, 0-24")	Midseason
	Leaf nutrients	Midseason
	Carbon fractions & enzyme analyses	Planting


Questions:




▲ New, Sac △ New, Woodland
● Old, Sac 1 ● Old, Sac 2 ○ Old, Woodland



“What factors look most different between old and new fields?”



“What factors positively correlate with yields, both between and within fields?”



“What factors start to look more alike over time?”

YIELDS & QUALITY




Soil type	Old/New	Variety	Harvest date	Yield (t/a)*	Brix	Yield % difference	Brix % difference
Tehama loam	Old	SVTM 9027	8-Aug	65.6	5.68	17	-7
	New	SVTM 9027	4-Aug	76.5	5.3		
Brentwood silty clay loam	Old	HM 58841	18-Aug	80.5	5.33	10	-10
	New	HM 58841	31-Jul	88.5	4.81		
Yolo silt loam	Old	HM 0371	1-Aug, 12-Aug	79.8	5.05	27	2
	New	HM 0371	30-Aug	101.6	5.13		





“WHAT LOOKS DIFFERENT?”

- Nutrition
- Compaction
- Disease



HIGH POTASSIUM IN ‘NEW’






**Yolo silt loam “Old”, pre-harvest
(8/1; 127 DAP)**

(Also consistently different between old & new: available phosphorus, available magnesium, leaf copper, leaf sodium)



**Yolo silt loam “New”, pre-harvest
(8/30; 148 DAP)**

SUBSURFACE COMPACTION









Brentwood silty clay loam “Old”

Bulk density=
1.52 g/cm³



Brentwood silty clay loam “New”

Bulk density=
1.34 g/cm³



Tehama New

Tehama Old

Massive
structure under
the drip tape.
Note roots
growing along
breakage face



topsoil







DISEASE

Hypothesis:

Sterilized soils will outyield non-sterilized soils in “old” fields, but not “new” ones






Fusarium stem rot & decline (FRD); F. foot rot




FRD; southern blight



No disease



Fusarium wilt (F2); southern blight





Rhizoctonia; charcoal rot



FRD; Verticillium wilt

WHAT CORRELATES TO YIELD?



w/in & between field effects of x are consistent
 ⇒ strong evidence for x forcing > yield

new2

old1 old2 new1 new2


field average

subsample or stratifying w/in field

possible variable causing > yield in new fields
 e.g. water content (15cm)

PROPOSED NEXT STEPS: 2026

- Measure change over time (2025 “New” fields)
- Measure “Old” vs “New” (from row crops)





TESTING A NEWLY REGISTERED NEMATICIDE : SALIBRO (CORTEVA)

7/20/23

SALIBRO FIELD TRIAL

- Field near Dixon, CA
- Resistance-breaking RKN-- yield loss est. ~8 t/a in 2023


2025: 5 replicates; every other row switched off


- Planted 4/21 (var HM 8237)
- Harvest 9/19

2025 Salibro program:

- 1st application—30.7 fl oz/acre, 20 DAP (1st irrigation after incorporation)
- 2nd application—30.7 fl oz/acre, 50 DAP
- At each application: mixed 30.7 fl oz/10 gal water, injected 10 gpa
- Cost about \$200/acre







- **11 t/a difference in trial rows**
- Difference in # dead plants pre-harvest not significant
- No difference in sunburn
- Yield difference likely due to delayed onset of symptoms



Acknowledgements

Hosting

- **Blake Harlan & Chris McAlister** (Harlan Family Ranch)
- **Bruce Rominger** (Rominger Brothers Farms)
- **Tim Beeman & Toshi Aoki** (Bullseye Farms)
- **Colin Muller** (M Three Ranches)
- **Spencer Bei & Aaron Black** (Robben Ranch)
- **Planting & harvest crews!!**

Donation of plants

- **Kevin Winn & Scott Picanso** (TS&L)

Field help, analytics & use of lab space

- **Jorge Rodrigues** (UC Davis Professor of Soil Ecology) & Lab
- **Daniel Geisseler** (UC Davis CE Specialist in Nutrient Management)
- **Sutie Xu** (UC Davis CE Specialist in Soil Health)
- **Cassandra Swett** (UC Davis CE Specialist in Plant Pathology)
- **Amanda Hodson** (UC Davis Professor of Nematology)
- **Shah Alam** (MS Student, UC Davis Dept of Nematology))
- **Eric Haddix** (UC Davis Vegetable Crops Facility)
- **Indiana Waterman, Mat Talton & UCCE colleagues** for harvest help (!!)

California Tomato Research Institute for funding

Corteva (Daniel Abruzzini) Salibro field day sponsorship



QUESTIONS?

PALAZICKI@UCANR.EDU; 530-219-5198